

Inference in Hidden Markov Models I: Local Asymptotic Normality in the Stationary Case

P. J. Bickel^{1 2}

University of California, Berkeley

and

Y. Ritov^{1 2 3}

Hebrew University, Jerusalem

Revision: April 1995

Abstract

Following up on Baum and Petrie (1966) we study likelihood based methods in hidden Markov models, where the hiding mechanism can lead to continuous observations and is itself governed by a parametric model. We show that procedures essentially equivalent to maximum likelihood estimates are asymptotically normal as expected and consistent estimates of their variance can be constructed, so that the usual inferential procedures are asymptotically valid.

AMS Classification: Primary: 62M09, Secondary: 62F12

Key words and phrases: maximum likelihood, hidden Markov models, local asymptotic normality, geometric ergodicity

¹Research partially supported by NSF Grant DMS-9115577

²Research partially supported by US/Israel Bi-National Science Foundation Grant ³ 90-00031/2

1 Introduction and basic results

Hidden Markov models, that is stochastic point functions of finite Markov chains, have become important in a number of areas of application. These include, first and foremost, speech recognition, see Rabiner (1989) for an introduction and survey, the study of excitation periods in ion channels, see Ball and Rice (1992) for a survey, models for heterogenous DNA sequences, Churchill (1992), among others. The main focus of these efforts have been algorithms for the fitting of these models and, in particular, see Rabiner, the implementation of likelihood based methods. It is, in fact, not obvious that the likelihood can be computed in linear time. But that is the case. There has been comparatively little work on the study of the inferential properties of likelihood methods in these models. The notable exceptions to this are the papers of Baum and Petrie (1966) and Petrie (1969) and most recently Leroux (1989), (1991). Concurrently with our work Rydén (1994a, 1994b) has also pursued likelihood based procedures in hidden Markov models.

Specifically, Baum and Petrie showed that, when observing a *deterministic finite point function* of a finite Markov chain, maximum likelihood estimates of the parameters of the model governing the chain are consistent and asymptotically normal. Leroux formulated hidden Markov models in the generality we shall present and established consistency of maximum likelihood estimates of both the parameters of the Markov chain and the conditional distribution of the observations given the Markov chain. Unlike the Baum-Petrie techniques, which were used both for establishing consistency and asymptotic normality, Leroux's approach based on results of Furstenberg and Kesten (1960) and Kingman's subadditive ergodic theorem (1976) appears incapable of giving results beyond consistency. On the other hand we shall show, by adding a few essential ideas to the penetrating analysis of Baum and Petrie, that the log likelihood for hidden Markov models obeys the local asymptotic normality (LAN) conditions of LeCam (see LeCam and Yang (1990), for instance). Hence, efficient analogues of maximum likelihood estimates can be constructed, and the information bound giving their asymptotic variance estimated. We shall also indicate how our results need to be strengthened to yield asymptotic efficiency of maximum likelihood estimates, when they are consistent. Consistency of maximum likelihood estimates can also be established with our methods but under conditions slightly stronger than those of Leroux (1991).

The paper is constructed as follows. In the rest of this section we formally introduce the models we consider, state our main assumptions and results, and further discuss the strengths and weaknesses of these as well as extensions and further questions, some of which we intend to pursue. In section 2 we give without proof some lemmas needed to establish our main theorem, discuss the heuristic behind them, and give a proof of the theorem based on these lemmas. Finally in section 3 we state more lemmas, give the proofs of all the lemmas which may not immediately be derived from the work of Baum and Petrie or others.

R1: Changed

Changed

Formally we assume that observations $(Y_1, \dots, Y_n) \in \mathcal{Y}^n$, for some space \mathcal{Y} , are distributed according to $P_\vartheta^{(n)}$, $\vartheta \in \Theta$ open $\subset R^p$ and described as follows:

- i) (Hidden chain) We are given (but do not observe) a stationary ergodic Markov chain X_1, \dots, X_n, \dots with states $\{1, \dots, K\}$, stationary initial probability $\pi_\vartheta(i)$, $1 \leq i \leq K$ and transition probability matrix $\|\alpha_\vartheta(i, j)\|_{K \times K}$.
- ii) (Y_i is a function of the present X_i and an external randomization only.) Given X_1, \dots, X_n the Y_i are conditionally independent, and given X_i , Y_i is independent of X_j , $j \neq i$.
- iii) (Stationarity) The conditional distribution of Y_i given X_i doesn't depend on i .
- iv) The conditional distributions of Y_i given $X_i = a$ are dominated by ν , a σ finite measure for all i, a, ϑ . The conditional density is denoted by $g_\vartheta(\cdot | a)$.

We may then write the density of (Y_1, \dots, Y_n) with respect to product measure $\nu^{(n)}$ as,

$$g_\vartheta(y_1, \dots, y_n) = \sum_{(x_1, \dots, x_n)} f_\vartheta(x_1, \dots, x_n, y_1, \dots, y_n) \quad (1.1)$$

where

$$f_\vartheta(x_1, \dots, x_n, y_1, \dots, y_n) = \pi_\vartheta(x_1) \prod_{j=1}^{n-1} \alpha_\vartheta(x_j, x_{j+1}) \prod_{i=1}^n g_\vartheta(y_i | x_i) \quad (1.2)$$

is the joint density of $(X_1, \dots, X_n, Y_1, \dots, Y_n)$ with respect to (counting measure) $^{(n)} \times \nu^{(n)}$. We denote the joint distribution of (X_i, Y_i) , $1 \leq i < \infty$, by P_ϑ , a probability on (Ω, \mathcal{A}) where Ω is the space of x, y sequences and \mathcal{A} is the Borel σ field.

This model, more or less given in Leroux (1989), is more general than it appears to be at first sight. It includes all situations where $Y_i = h(X_{i-j}; 1 \leq j \leq t, \epsilon_i, \vartheta)$, $1 \leq i \leq n$ where the ϵ_i are i.i.d. and independent of the X 's and t is fixed, since we can always take $(X_{1+i}, \dots, X_{t+i})$ $i \geq 0$ as our hidden chain. We will need the following assumptions.

A1: For all ϑ, a, b , $\alpha_\vartheta(a, b) \geq \gamma(\vartheta) > 0$.

A2: For all a, b , the map $\vartheta \rightarrow \alpha_\vartheta(a, b)$ has three continuous derivatives. Hence so has $\vartheta \rightarrow \pi_\vartheta(a)$.

R2: Changed
Was it needed?

Note that A1 and A2 imply that for all ϑ_0 there exists $\delta > 0$, $\gamma(\vartheta_0) > 0$ such that

$$\inf\{\alpha_\vartheta(a, b) : |\vartheta - \vartheta_0| \leq \delta\} \geq \gamma(\vartheta_0) \quad (1.3)$$

$$\inf\{\pi_\vartheta(a) : |\vartheta - \vartheta_0| \leq \delta\} \geq \gamma(\vartheta_0). \quad (1.4)$$

A3: The maps $\vartheta \rightarrow \nabla \log g_\vartheta(y|a)$ has three derviatives for all y, a . Further, for all ϑ_0 there exists $\delta > 0$, $\lambda > 0$ such that if,

R3: Changed

$$q_{\vartheta_0}(y, \delta) \equiv \sup\{|\nabla \log g_\vartheta(y|a)| : a, |\vartheta - \vartheta_0| \leq \delta\}$$

then

$$E_{\vartheta_0} \exp[\lambda q_{\vartheta_0}(Y_1, \delta)] < \infty \quad (1.5)$$

A4: For all ϑ_0 there exists $\delta > 0$, $r > 32$ such that if,

It was $r > 16$.

$$\rho_{\vartheta_0}(y) = \sup\left\{\frac{g_\vartheta(y|a)}{g_\vartheta(y|b)} : a, b, |\vartheta - \vartheta_0| < \delta\right\}$$

then,

$$E_{\vartheta_0} \rho_{\vartheta_0}^r(Y_1) < \infty \quad (1.6)$$

A5: Let $\vartheta = (\vartheta_1, \dots, \vartheta_p)$ and

$$q_{\vartheta_0 j}(y, \delta) = \sup\left\{\left|\frac{\partial^j}{\partial \vartheta_{i_1} \dots \partial \vartheta_{i_j}} \log g_{\vartheta}(y|a)\right|\right\},$$

where the sup is taken over $\{1 \leq i_l \leq p, l = 1, \dots, j, 1 \leq a \leq K, |\vartheta - \vartheta_0| \leq \delta\}$. Assume for all ϑ_0 , some $\delta > 0$, $j = 2, 3$

$$E_{\vartheta_0} \left\{ (q_{\vartheta_0 j}(Y_1, \delta))^{2+\delta} \right\} < \infty. \quad (1.7)$$

Let (X_i, Y_i) , $-\infty < i < \infty$ be the two sided stationary sequence defined by our model and,

$$W(Y_1, Y_0, Y_{-1}, \dots) \equiv \sum_{m=-\infty}^1 W_m(Y_1, Y_0, \dots) \quad (1.8)$$

R4: Changed
All the
terms for $m =$
1 were added,
most are 0.

where

$$\begin{aligned} W_m(Y_1, Y_0, \dots) & \quad (1.9) \\ \equiv & E_{\vartheta_0} \{ \nabla \log g(Y_m | X_m) | Y_1, Y_0, \dots \} - E_{\vartheta_0} \{ \nabla \log g(Y_m | X_m) | Y_0, Y_{-1}, \dots \} \\ & + E_{\vartheta_0} \{ \nabla \log \alpha(X_m, X_{m+1}) | Y_1, Y_0, \dots \} \\ & - E_{\vartheta_0} \{ \nabla \log \alpha(X_m, X_{m+1}) | Y_0, Y_{-1}, \dots \} \end{aligned}$$

We show in Lemma 3.5 that, under A1-A4, $W \in L_2(P_{\vartheta_0})$ and we can then define,

$$I(\vartheta_0) \equiv E_{\vartheta_0} \{ W W^T \}. \quad (1.10)$$

Fix ϑ_0 and let \mathcal{L}_0, P_0, E_0 be law, probability and expectation under ϑ_0 . Let $\delta_n \equiv n^{-1/2}$, $\vartheta_n \equiv \vartheta_0 + \tau \delta_n$, and

$$L_n(\tau) \equiv \frac{g_{\vartheta_n}}{g_{\vartheta_0}}(Y_1, \dots, Y_n). \quad (1.11)$$

Our main goal is to establish the following,

Theorem 1.1 *Suppose assumptions A1-5 hold. Then there exist Δ_n , random p vectors, such that, if $|\tau_n| = O(1)$,*

R5: Changed

$$\log L_n(\tau_n) = \tau_n^T \Delta_n - \frac{1}{2} \tau_n^T J_n \tau_n + R_n(\tau_n) \quad (1.12)$$

where

$$E_0 \Delta_n = 0, \quad (1.13)$$

$$E_0 \Delta_n \Delta_n^T \rightarrow I(\vartheta_0), \quad (1.14)$$

$$J_n \rightarrow I(\vartheta_0), \quad (1.15)$$

$$\Delta_n \xrightarrow{\mathcal{L}_0} \mathcal{N}(0, I(\vartheta_0)), \quad (1.16)$$

$P_0(|R_n(\tau_n)| > n^{-\gamma/2}/e_n) < \max\{e_n, n^{-1}\}$ for any $e_n \rightarrow 0$ and $\gamma < 2(1 - 16/r)/5$ for r satisfying (1.6), and $I(\vartheta_0)$ given in (1.10).

Note that (1.12) is just local asymptotic normality (LAN) in the sense of LeCam. In order to implement this result for inferential purposes we can proceed more or less as in LeCam and Yang (1990) pp. 57-65. We need

A6: The parameter ϑ is identifiable in the sense that if for some $\vartheta, \vartheta' \in \Theta$
 $P_{\vartheta}^{(n)} = P_{\vartheta'}^{(n)}$ for all n , then $\vartheta = \vartheta'$.

R6: Not
changed

Lemma 1.1 *If A1–A6 hold, then there exists an estimate $\{\tilde{\vartheta}_n(Y_1, \dots, Y_n)\}_{n \geq 1}$ which is \sqrt{n} consistent. That is, for each ϑ_0 , $\tilde{\vartheta}_n - \vartheta_0 = O_{P_0}(\delta_n)$.*

Let the \mathcal{G}_n grid denote the set of all $(\pm j_1, \dots, \pm j_p) \delta_n n^{-\gamma/2p}$ where the j_i are integers and γ is as in Theorem 1.1. If Lemma 1.1 holds we can and shall without loss of generality suppose that $\tilde{\vartheta}_n$ takes on values in the \mathcal{G}_n grid only. Let,

$$\hat{\vartheta}_n = \text{local maximizer of } g_{\vartheta}(Y_1, \dots, Y_n) \text{ on } \mathcal{G}_n \quad (1.17)$$

closest to $\tilde{\vartheta}_n$ among the points of the δ_n grid, and for given ϵ_n , define the matrix \hat{I}_n by,

R7: Changed

$$\hat{I}_{nab} = -\epsilon_n^{-2} \log \left\{ \frac{g_{\hat{\vartheta}_n(a,b)} g_{\hat{\vartheta}_n} (Y_1, \dots, Y_n)}{g_{\hat{\vartheta}_n(a,0)} g_{\hat{\vartheta}_n(0,b)}} \right\} \quad (1.18)$$

$$\hat{\vartheta}_n(a, b) = \hat{\vartheta}_n + \epsilon_n \delta_n (e_a + e_b) \quad (1.19)$$

where e_1, \dots, e_p are the standard basis vectors and $e_0 = 0$. Thus, $\hat{\vartheta}_n$ is a grid version of the closest root of the likelihood equation to $\tilde{\vartheta}_n$ and $-\hat{I}_n$ is a second difference grid evaluated version of the Hessian at $\hat{\vartheta}_n$. Then,

R8: Changed

Corollary 1.1 *If A1 - A6 hold, $\hat{\vartheta}_n$ is as in (1.17) and $I(\vartheta_0)$ is nonsingular, then,*

R9: Changed

$$n^{1/2}(\hat{\vartheta}_n - \vartheta_0) \xrightarrow{\mathcal{L}_0} \mathcal{N}(0, I^{-1}(\vartheta_0)) \quad (1.20)$$

$$\hat{I}_n \xrightarrow{P_0} I(\vartheta_0). \quad (1.21)$$

We are now able to construct asymptotically efficient estimates, tests, etc by pretending that $\hat{\vartheta}_n$ is approximately $\mathcal{N}(\vartheta, \delta_n^2 \hat{I}^{-1})$. This result does not give what one would ideally like:

- a) That the M.L.E. $\hat{\vartheta}_n^*$ is asymptotically normal $(\vartheta_0, \delta_n^2 I^{-1}(\vartheta_0))$
- b) That the Hessian of the log likelihood at $\hat{\vartheta}_n^*$, $n^{-1} \left\| \frac{\partial^2}{\partial \vartheta_a \partial \vartheta_b} \log g_{\hat{\vartheta}_n^*}(Y_1, \dots, Y_n) \right\|$ converges in probability to $-I(\vartheta_0)$.

Part a) requires \sqrt{n} consistency of the MLE and uniform (permitting τ_n to be data determined) LAN while b) requires consistency of the MLE and some sort of uniform convergence of the Hessian. These are open problems.

Discussion of assumptions:

Evidently using f_{ϑ} and Bayes rule we can construct maps from \mathcal{Y}^n to $\{\text{Probabilities on } (\Omega, \mathcal{A})\}$, $(y_1, \dots, y_n) \rightarrow P_{\vartheta}(\cdot | y_1, \dots, y_n)$ such that $P_{\vartheta}(\cdot | Y_1, \dots, Y_n)$ is a regular conditional probability on Ω given (Y_1, \dots, Y_n) . The key property in Baum and Petrie's and our analysis is that (X_1, X_2, \dots) are an inhomogeneous Markov chain under $P_{\vartheta}(\cdot | y_1, y_2, \dots)$. Assumptions A1, A2, and A4 guarantee that, with probability 1, this chain has strong geometric ergodicity properties which among other things guarantee the existence of $I(\vartheta_0)$ in (1.10). A1 and A2 can easily be relaxed by specifying that only some power of the transition matrix needs to have all entries positive. A4 is clearly not very demanding. A3 intersects with A1, A2, and A4 guaranteeing the validity of appropriate Taylor expansions. It is evidently a much stronger moment condition than what is required for valid Taylor expansions in the i.i.d. case. However, we do not presently see how it can be relaxed. It evidently holds for Gaussian location and scale families, for instance, as does A5 which is essentially a standard condition of the Cramér type.

Extensions: Two extensions worth considering are,

- a) To drop the requirement that the state space of X be finite.

b) To the case where the hidden process is a Markov random field.

The first extension includes most nonlinear ARMA processes which have been proposed – see Priestley (1988), Tong (1991). Let $\dots, \epsilon_{-1}, \epsilon_0, \epsilon_1, \dots$ be an iid sequence of random variables with distribution from a parametric family, $\{F_\vartheta\}$, and

$$Y_j = h(\epsilon_j, \epsilon_{j-1}, \dots, \vartheta), \quad 1 \leq j \leq n. \quad (1.22)$$

Since $X_j = \{\epsilon_{j-k} : k \geq 0\}$ is a Markov chain on R^∞ this falls under case a). For a discussion of Edgeworth expansions of smooth statistics in such models see Götze and Hipp (1992).

Estimation of parameters in hidden Markov fields by ad hoc methods has been considered by Frigessi (1990) and others. Likelihoods even for directly observed fields are only computable by simulation but extension of our approach replacing likelihoods of the hidden process by pseudo likelihoods may be valuable. See Qian and Titterton (1991)

We intend to pursue special cases of both extensions. It also appears that extensions to continuous time situations where observations are not simply point functions of the hidden process may also be possible and interesting. A simple example discussed in Daley and Vere Jones (1989), and pursued by Rydén (1994b), is that of Cox processes driven by a finite state continuous Markov process.

R10: Changed

2 Proof of theorem 1.1

We begin with an outline of our proof of theorem 1.1. Details are given at the end of the section after the statement of some lemmas. Let $\mathbf{Y}_{a,b} = (Y_a, \dots, Y_b)$ and $\mathbf{X}_{a,b}$ be the corresponding X block. Also define $\mathbf{Y}_m^{(k)} = \mathbf{Y}_{mk+1, mk+k}$ and $\mathbf{X}_m^{(k)}$ be the corresponding X block where $0 \leq m \leq N = n/k - 1$. To simplify the notation we assume that n is a multiple of k . We argue in II below that if k does not divide n we can neglect the resulting end effect. For convenience we use the subscript τ in the sequel to stand for $\vartheta_n = \vartheta_0 + \tau_n \delta_n$, where $\{\tau_n\}$ is a bounded sequence. Let, $\ell_\tau(\mathbf{Y}_m^{(k)} | X_{mk+1})$

denote the conditional likelihood of $\mathbf{Y}_m^{(k)}$ given X_{mk+1} and let

$$L_{\tau m} \equiv \frac{\sum_{a=1}^K P_\tau[X_{mk+1} = a | \mathbf{Y}_{1,mk}] \ell_\tau(\mathbf{Y}_m^{(k)} | a)}{\sum_{a=1}^K P_0[X_{mk+1} = a | \mathbf{Y}_{1,mk}] \ell_0(\mathbf{Y}_m^{(k)} | a)}, \quad (2.1)$$

denote the likelihood ratio of $\mathbf{Y}_m^{(k)}$ given $\mathbf{Y}_{1,mk}$. Also, let

$$L_{\tau m}^{(d)} \equiv \frac{\sum_{a=1}^K P_\tau[X_{mk+1} = a | \mathbf{Y}_{mk-d,mk}] \ell_\tau(\mathbf{Y}_m^{(k)} | a)}{\sum_{a=1}^K P_0[X_{mk+1} = a | \mathbf{Y}_{mk-d,mk}] \ell_0(\mathbf{Y}_m^{(k)} | a)}, \quad (2.2)$$

denote the likelihood ratio of $\mathbf{Y}_m^{(k)}$ given $\mathbf{Y}_{mk-d,mk}$, and

$$L_{\tau m}^* \equiv \frac{\ell_\tau(\mathbf{Y}_m^{(k)} | X_{mk+1})}{\ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1})} \quad (2.3)$$

the likelihood ratio of $\mathbf{Y}_m^{(k)}$ given X_{mk+1} .

I. Write

$$\log L_n(\tau) = \sum_{m=1}^N \log L_{\tau m} + \log \frac{g_{\partial n}}{g_{\partial_0}}(Y_1, \dots, Y_k) \quad (2.4)$$

and

$$\sum_{m=1}^N \log L_{\tau m} = \sum_{m=1}^N \log L_{\tau m}^* + \sum_{m=1}^N \log \left(1 + \frac{(L_{\tau m} - L_{\tau m}^*)}{L_{\tau m}^*} \right). \quad (2.5)$$

Taylor expanding we get

$$\begin{aligned} & \sum_{m=1}^N \log \left(1 + \frac{(L_{\tau m} - L_{\tau m}^*)}{L_{\tau m}^*} \right) \\ &= \sum_{m=1}^N (L_{\tau m} - L_{\tau m}^*) - \sum_{m=1}^N \frac{(L_{\tau m} - L_{\tau m}^*)}{L_{\tau m}^*} (L_{\tau m}^* - 1) \\ & \quad - \frac{1}{2} (1 + R_n) \sum_{m=1}^N \frac{(L_{\tau m} - L_{\tau m}^*)^2}{(L_{\tau m}^*)^2}. \end{aligned} \quad (2.6)$$

II. We expect $|L_{\tau m} - 1| = O_{P_0}(k/n)^{1/2}$. We shall establish this and in so doing also show that if $n = Nk + r$, $0 < r < k$, then the difference between

$\log L_n(\tau)$ and $\log L_{Nk}(\tau)$ is $o_{P_0}(1)$. Further, X_1, X_2, \dots remains a Markov chain given the Y 's. Although the chain is not stationary, it satisfies a strong mixing condition. Thus, we expect that the knowledge of Y 's and X 's in the distant past adds very little information to the present and $|L_{\tau m} - L_{\tau m}^*| = o_{P_0}((k/n)^{1/2})$ so that we can and do show that the last two terms of (2.6) are negligible. The second term in (2.4) is also negligible. This uses arguments based on the Baum-Petrie results which are stated under our conditions in lemmas 3.1–3.4. R11: Changed
R12: Changed

III. We write the first term as

$$\sum_{m=1}^N (L_{\tau m} - L_{\tau m}^*) = \sum_{m=1}^N (L_{\tau m}^{(d)} - L_{\tau m}^*) + \sum_{m=1}^N (L_{\tau m} - L_{\tau m}^{(d)}). \quad (2.7)$$

We show that the second term is negligible for $d \rightarrow \infty$, $d = o(k)$ using Baum-Petrie again and that the first term is negligible using uniform mixing and the Ibragimov-Linnik lemma (Lemma 3.7 below). R13: Changed

IV. We Taylor expand $\sum_{m=1}^N \log L_{\tau m}^*$ in τ and apply uniform mixing to show it has the LAN structure.

Finally,

V. We evaluate $I(\vartheta_0)$ necessarily by a different starting formula than Baum-Petrie's, but again rely on their results to dispose of possible long range dependence.

The proof of Theorem 1.1 is based on the following lemmas whose proofs are given in the next section.

We adopt the following notation. We say

$$A_n = O_{b_n}(a_n) \quad (2.8)$$

iff there exists some $M_0, c(\cdot) \searrow 0$ such that for all $M > M_0$ and $n > n(M)$

$$P_0[|A_n| \geq M a_n] \leq c(M) b_n.$$

In particular, $O_0(a_n) \equiv O(a_n)$ and $O_1(a_n) \equiv O_{P_0}(a_n)$.

Lemma 2.1 *If A1 - A4 hold, $r > 16$, $k = n^{4\epsilon + \frac{5}{4}\gamma}$, $\epsilon > \frac{2}{r}$, $4\epsilon + \gamma < \frac{1}{2}$, $\gamma > 0$ then for any $|\tau| < M$,* R14: Changed

$$\sum_{m=1}^N \log(L_{\tau m}/L_{\tau m}^*) = O_{e_n}(n^{-\gamma/2}/e_n) \quad (2.9)$$

for any $e_n \rightarrow 0$, $n e_n \rightarrow \infty$

Lemma 2.2 *If A1 - A5 hold, $r > 32$, $k = n^{4\epsilon + \gamma}$, $4\epsilon + \gamma < \frac{1}{4}$ then,*

$$E_0 \sup_{|\tau| < M} \left\{ \sum_{m=1}^N \left| \log L_{\tau m}^* - \delta_n \tau^T \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \right. \right. \\ \left. \left. - \frac{1}{2n} \tau^T \left\| \frac{\partial^2}{\partial \vartheta_i \partial \vartheta_j} \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \right\| \tau \right\} \\ = O(k^2/n^{1/2}), \quad (2.10)$$

The lemma is now weaker ($k = o(n^{-1/4})$) and not $k = o(n^{-1/2})$.

where $\|a_{ij}\|$ is the matrix with entries a_{ij} .

Lemma 2.3 *Under A1 - A4*

$$\lim_{k \rightarrow \infty} \frac{1}{k} E_0 \left\{ \left(\nabla \log \ell_0(\mathbf{Y}_0^{(k)} | X_1) \right) \left(\nabla \log \ell_0(\mathbf{Y}_0^{(k)} | X_1) \right)^T \right\} = I(\vartheta_0) \quad (2.11)$$

where $I(\vartheta_0)$ is defined as in (1.10).

Lemma 2.4 *Under A1 - A4, if $k = o(n)$*

$$\frac{1}{n} \sum_{m=1}^N E_0 \{ \nabla \nabla^T \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) | X_{mk+1} \} \xrightarrow{P_0} I(\vartheta_0) \quad (2.12)$$

$$\frac{1}{n} \sum_{m=1}^N E_0^{1/2} \{ | \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) |^4 | X_{mk+1} \} = O_{P_0}(1) \quad (2.13)$$

$$\max_m P_0 [| \delta_n \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) | \geq \epsilon | X_{mk+1}] = o_{P_0}(1) \quad (2.14)$$

where $\nabla \nabla^T h \equiv (\nabla h)(\nabla h)^T$.

Lemma 2.5 *Under A1 - A4,*

$$\frac{1}{n} \sum_{m=1}^N \left\| \frac{\partial^2}{\partial \vartheta_a \partial \vartheta_b} \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \right\| \xrightarrow{P_0} -I(\vartheta_0) \quad (2.15)$$

Proof of Theorem 1.1: From lemma 2.1 we see that if $\tau \equiv \tau_n$ we can replace the left hand side of (2.5) by $\sum_{m=1}^N \log L_{\tau_n m}^* + O_{\epsilon_n} \left(n^{-2\gamma/5} / \epsilon_n \right)$ if

R15: Changed Since, γ is now smaller we have no problem. The last line of the display was changed, since Lemma 2.2 is weaker.

$k = n^{4\epsilon + \gamma}$, $\epsilon > \frac{2}{r}$, $4\epsilon + \gamma < \frac{1}{4}$.

Lemma 2.2 now guarantees that

$$\begin{aligned} \sum_{m=1}^N \log L_{\tau_n m}^* &= \delta_n \tau_n^T \sum_{m=1}^N \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \\ &+ \frac{1}{2n} \sum_{m=1}^N \tau_n^T \left\| \frac{\partial^2 \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1})}{\partial \vartheta_i \partial \vartheta_j} \right\| \tau_n \\ &+ O_{\epsilon_n} \left(n^{-1/2 + 8\epsilon + 2\gamma} / e_n \right). \end{aligned} \quad (\text{a})$$

Let

$$\xi_{mn} = \delta_n \tau_n^T \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}), \quad 1 \leq m \leq N. \quad (\text{b})$$

We claim that this is a triangular sequence of martingale summands with respect to the σ fields $\mathcal{F}_{mn} = \sigma(\mathbf{X}_{1, (m+1)k+1}, \mathbf{Y}_{1, (m+1)k})$, $1 \leq m \leq N$. This follows from the Markov property which gives,

$$E_0 \left\{ \frac{\ell_{\vartheta}}{\ell_0}(\mathbf{Y}_m^{(k)} | X_{mk+1}) | \mathcal{F}_{(m-1)n} \right\} = E_0 \left\{ \frac{\ell_{\vartheta}}{\ell_0}(\mathbf{Y}_m^{(k)} | X_{mk+1}) | X_{mk+1} \right\} \equiv 1 \quad (\text{c})$$

and the usual interchange of differentiation and integration. Further, $I(\vartheta_0)$ is well defined and by Lemma 2.4 equation (2.12),

R16: Changed

$$\sum_{m=1}^N E_0(\xi_{mn}^2 | \mathcal{F}_{(m-1)n}) \xrightarrow{P_0} \tau_n^T I(\vartheta_0) \tau_n, \quad (\text{d})$$

and by lemma 2.4,

$$\begin{aligned} &\sum_{m=1}^N E_0(\xi_{mn}^2 \mathbf{1}(|\xi_{mn}| \geq \epsilon) | \mathcal{F}_{(m-1)n}) \\ &\leq \left[\sum_{m=1}^N E_0^{1/2}(\xi_{mn}^4 | \mathcal{F}_{(m-1)n}) \right] \max_{1 \leq m \leq N} P_0^{1/2}[|\xi_{mn}| \geq \epsilon | \mathcal{F}_{(m-1)n}] \\ &= o_{P_0}(1) \end{aligned} \quad (\text{e})$$

The central limit theorem for triangular arrays of martingale summands (see Hall and Heyde (1980) for example) establishes that

R17: Changed

$$\delta_n \tau^T \sum_{m=1}^N \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \xrightarrow{\mathcal{L}_0} \mathcal{N}(0, \tau^T I(\vartheta_0) \tau). \quad (\text{f})$$

Finally, lemma 2.5 establishes that the last term in (a) tends to $-\frac{1}{2}\tau^T I(\vartheta_0)\tau$. The theorem is proved. \square

Proof of Lemma 1.1: We construct a minimum distance estimator. The proof is based on LeCam (1956). The construction is simple under the assumption that for some $k < \infty$, the map $\vartheta \rightarrow P_\vartheta^{(k)}$ is 1-1 and Θ compact. In that case it is possible to construct \sqrt{n} consistent estimates by considering $P_n^{(k)}$, the empirical distribution of the vectors $\{Y_{a+b} : 0 \leq b \leq k-1\}$, for $1 \leq a \leq n-k+1$. See Rydén (1995) for a proof that $k = 2K$ under some what different conditions than ours and Rydén (1994a) for the construction of the \sqrt{n} consistent estimator. In general, let $\Theta = \bigcup_{j=1}^\infty \Theta_j$ with $\Theta_{j+1} \supset \Theta_j$, $j \geq 1$ compact sets, and define $T_{njk} = \{t \in \Theta_j : n^{-1/4}d_K(P_t^{(k)}, P_n^{(k)}) = \min_{\vartheta \in \Theta_j} d_K(P_\vartheta^{(k)}, P_n^{(k)})\}$ where $d_K(\cdot, \cdot)$ is the Kolmogorov distance. Then let $\tilde{\vartheta} \in T_n$ where $T_n = T_{njk}$ with T_{njk} non-empty and radius less than $n^{-1/4}$ and minimal $j+k$. \square

Proof of Corollary 1.1: The corollary follows in a standard fashion by the methods of LeCam (1986) and LeCam and Yang (1990). Let $\mathcal{G}_{Mn} = \mathcal{G}_n \cap \{\vartheta : |\vartheta - \vartheta_0| < Mn^{-1/2}\}$. Note that there are $O(n^{\gamma/2})$ points in \mathcal{G}_{Mn} . Write $R_n = R_n(\tau)$ for the remainder term in (1.12). It follows from Theorem 1.1 that

$$P_0 \left(\sup_{\tau n^{-1/2} \in \mathcal{G}_{Mn}} |L_n(\tau) - \tau \Delta_n + 1/2\tau^T J_n \tau| > \epsilon \right) \quad (\text{a})$$

$$\leq O(n^{\gamma/2}) \sup_{\tau n^{-1/2} \in \mathcal{G}_{Mn}} P_0(|R_n(\tau)| > \epsilon) \quad (\text{b})$$

$$\xrightarrow{P_0} 0.$$

Hence $\hat{\vartheta}_n$ is in distance $O_{P_0}(n^{(1+\gamma)/2})$ of

$$n^{-1/2} \arg \max \{ \tau^T \Delta_n - 1/2\tau^T J_n \tau \} = n^{-1/2} J_n^{-1} \Delta_n, \quad (\text{c})$$

which proves Corollary 1.1. \square

3 Further lemmas and proofs

We begin with four lemmas which are straightforward extensions of key results of Baum and Petrie (1966) (lemma 2.1, lemma 2.2, and corollary 2.3) valid under assumptions A1 - A3 and hence the proofs are omitted. They contain the essential information that knowledge of y 's and x 's in the distant past adds very little information to the present. Lemma 3.1 guarantees strong mixing conditions.

Let

$$\mu_0(y) = \left(1 + (K - 1)\gamma^{-2}(\vartheta_0)\rho_{\vartheta_0}(y)\right)^{-1}$$

In what follows we write $P_\vartheta(A|B, y_1, \dots, y_n)$ if $P_\vartheta(A|B, Y_1, \dots, Y_n)$ is a version of the regular conditional probability of A given B, Y_1, \dots, Y_n , and $P_\vartheta(A|B, y_1, \dots, y_n)$ is defined for all ϑ, A, B and y_1, \dots, y_n . This is easily done if we can define densities $g_\vartheta(y|x)$ valid for all ϑ, y and x .

Lemma 3.1 For $|\vartheta - \vartheta_0| \leq \delta$ and all ϑ_0 ,

$$P_\vartheta[X_{i+1} = b|X_i = a, y_1, \dots, y_n] \geq \mu_0(y_{i+1}) > 0. \quad (3.1)$$

Lemma 3.2 If C_t is an event depending on $X_i, Y_i, i \geq t$, only, then for all $|\vartheta - \vartheta_0| \leq \delta, \vartheta_0, d \geq 2$,

$$\begin{aligned} |P_\vartheta[C_t|y_{t-1}, \dots, y_{t-d+1}] - P_{\vartheta_0}[C_t|y_{t-1}, \dots, y_{t-d}]| &\leq \prod_{j=t-d+1}^{t-1} (1 - 2\mu_0(y_j)) \\ &\leq \exp\left\{-2 \sum_{j=t-d+1}^{t-1} \mu_0(y_j)\right\} \end{aligned}$$

Lemma 3.3 Let C_t be as above and

$$M_d^+(\vartheta) = \max_a P_\vartheta[C_t|y_1, \dots, y_n, X_{t-d} = a]$$

and define $M_d^-(\vartheta)$ as the corresponding minimum. Then, for all $\vartheta_0, |\vartheta - \vartheta_0| \leq \delta$,

$$\left|M_d^+(\vartheta) - M_d^-(\vartheta)\right| \leq \prod_{j=t-d+1}^{t-1} (1 - 2\mu_0(y_j)) \quad (3.2)$$

Lemma 3.4 *If A1 and A2 hold then for all ϑ_0 , $|\vartheta - \vartheta_0| \leq \delta$, y_1, \dots, y_ℓ , a, b*

$$P_\vartheta[X_{\ell+1} = a | y_1, \dots, y_\ell, X_1 = b] \geq \gamma(\vartheta_0). \quad (3.3)$$

The following two lemmas are of general utility in missing data models.

Lemma 3.5 *If $P \gg Q$, $e^\Lambda \equiv \frac{dQ}{dP}$, $T \in L_1(Q)$, and \mathbf{B} is a sub σ -field, then*

$$E_P |E_Q(T|\mathbf{B})| \leq E_P^{\frac{1}{r}} \{|T|^r\} E_P^{\frac{1}{s}} \{e^{s\Lambda}\} E_P^{\frac{1}{t}} \{e^{-t\Lambda}\}, \quad (3.4)$$

where $\frac{1}{r} + \frac{1}{s} + \frac{1}{t} = 1$.

Proof of lemma 3.5: Note that,

$$E_Q(T|\mathbf{B}) = \frac{E_P(Te^\Lambda|\mathbf{B})}{E_P(e^\Lambda|\mathbf{B})} \quad (a)$$

So, (3.4) is bounded by

$$\begin{aligned} E_P |E_P(Te^\Lambda|\mathbf{B})E_P(e^{-\Lambda}|\mathbf{B})| &\leq E_P \left\{ |T|e^\Lambda E_P(e^{-\Lambda}|\mathbf{B}) \right\} \\ &\leq E_P^{\frac{1}{r}} \{|T|^r\} E_P^{\frac{1}{s}} \{e^{s\Lambda}\} E_P^{\frac{1}{t}} \{e^{-t\Lambda}\}. \end{aligned} \quad (b)$$

□

Lemma 3.6 *Let $\vartheta \rightarrow U_\vartheta$, $\vartheta \in R$, be continuously differentiable where $U_\vartheta(\cdot)$ is a stochastic process on (Ω, \mathcal{A}) , \mathbf{B} is a sub field of \mathcal{A} . Then, if $P_\vartheta \ll \nu$ and $\ell_\vartheta \equiv dP_\vartheta/d\nu$, suppose*

- (i) $\vartheta \rightarrow \frac{\partial}{\partial \vartheta} \log \ell_\vartheta$
- (ii) $\vartheta \rightarrow E_\vartheta \left| \frac{\partial U_\vartheta}{\partial \vartheta} \right|$
- (iii) $\vartheta \rightarrow E_\vartheta [U_\vartheta^2]$
- (iv) $\vartheta \rightarrow E_\vartheta \left(\frac{\partial}{\partial \vartheta} \log \ell_\vartheta \right)^2$

are all continuous.

Then,

$$\frac{\partial}{\partial \vartheta} E_\vartheta \left(U_\vartheta | \mathbf{B} \right) = E_\vartheta \left(\frac{\partial U_\vartheta}{\partial \vartheta} | \mathbf{B} \right) + \text{cov}_\vartheta \left\{ \left(U_\vartheta, \frac{\partial}{\partial \vartheta} \log \ell_\vartheta \right) | \mathbf{B} \right\} \quad (3.5)$$

Proof of lemma 3.6: Write, $\Lambda(\vartheta, \vartheta + \Delta) = \log(\ell_{\vartheta+\Delta}/\ell_{\vartheta})$,

$$E_{\vartheta+\Delta}(U_{\vartheta+\Delta}|\mathbf{B}) = \frac{E_{\vartheta}(U_{\vartheta+\Delta} e^{\Lambda(\vartheta, \vartheta+\Delta)}|\mathbf{B})}{E_{\vartheta}(e^{\Lambda(\vartheta, \vartheta+\Delta)}|\mathbf{B})}. \quad (\text{a})$$

Then

$$\begin{aligned} \frac{\partial}{\partial \vartheta} E_{\vartheta}(U_{\vartheta}|\mathbf{B}) &= \frac{\partial}{\partial \Delta} E_{\vartheta}(U_{\vartheta+\Delta} e^{\Lambda(\vartheta, \vartheta+\Delta)}|\mathbf{B})\Big|_{\Delta=0} \\ &\quad - E_{\vartheta}(U_{\vartheta}|\mathbf{B}) \frac{\partial}{\partial \Delta} E_{\vartheta}(e^{\Lambda(\vartheta, \vartheta+\Delta)}|\mathbf{B})\Big|_{\Delta=0} \end{aligned} \quad (\text{b})$$

provided the right hand side exists. Interchange of integration and differentiation may be justified under our condition by a delicate but standard argument we do not reproduce. We get that the right hand side of (b) is,

$$E_{\vartheta}\left(\frac{\partial U_{\vartheta}}{\partial \vartheta}\Big|\mathbf{B}\right) + E_{\vartheta}\left(U_{\vartheta} \frac{\partial}{\partial \vartheta} \log \ell_{\vartheta}\Big|\mathbf{B}\right) - E_{\vartheta}(U_{\vartheta}|\mathbf{B}) E_{\vartheta}\left(\frac{\partial}{\partial \vartheta} \log \ell_{\vartheta}\Big|\mathbf{B}\right) \quad (\text{c})$$

and (3.5) follows. □

We also need a basic lemma from Ibragimov and Linnik (1971) (theorem 17.2.2 (p. 307)) which we quote for completeness.

Lemma 3.7 *If ξ, η have joint distribution P with marginals P_1, P_2 such that $\|P - (P_1 \times P_2)\| \leq \alpha$ where $\|\cdot\|$ is variational distance and for some $\delta > 0$, and $E|\xi|^{2+\delta} \leq c_1, E|\eta|^{2+\delta} \leq c_2$ then*

$$|E(\xi\eta) - E(\xi)E(\eta)| \leq c\alpha^{1-\beta} \quad (3.6)$$

where $\beta = 2/(2 + \delta)$ and $c = 4 + 3c_1^{\beta/2}c_2^{1-\beta/2} + 3c_1^{1-\beta/2}c_2^{\beta/2}$.

Here are the additional lemmas we need to carry out I-V. Let

$$\alpha_{\tau, i, m}(a, b) \equiv P_{\tau}[X_{i+1} = b \mid X_i = a, Y_1, \dots, Y_m] \quad (3.7)$$

Lemma 3.8 *In our model, if $1 \leq i \leq m - 1$,*

$$\frac{\alpha_{\tau,i,m}(a,b)}{\alpha_{0,i,m}(a,b)} = \frac{E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) | X_i = a, X_{i+1} = b, \mathbf{Y}_{1,m} \right\}}{E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) | X_i = a, \mathbf{Y}_{1,m} \right\}}. \quad (3.8)$$

Proof of lemma 3.8: Note that,

$$\begin{aligned} P_\tau[X_{i+1} = b, X_i = a | \mathbf{Y}_{1,m}] & \quad (a) \\ &= \frac{E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) 1(X_{i+1} = b, X_i = a) | \mathbf{Y}_{1,m} \right\}}{E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) | \mathbf{Y}_{1,m} \right\}} \end{aligned}$$

$$P_\tau[X_i = a | \mathbf{Y}_{1,m}] = \frac{E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) 1(X_i = a) | \mathbf{Y}_{1,m} \right\}}{E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) | \mathbf{Y}_{1,m} \right\}} \quad (b)$$

$$\begin{aligned} & E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) 1(X_i = a) | \mathbf{Y}_{1,m} \right\} \quad (c) \\ &= E_0 \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{1,m}, \mathbf{Y}_{1,m}) | X_i = a, \mathbf{Y}_{1,m} \right\} P_0[X_i = a | \mathbf{Y}_{1,m}] \end{aligned}$$

Substitute (a), (b) on the LHS of (3.8) and simplify using (c) and an analogous expression for the numerator in (a) to get the RHS.

□

Let

$$S_n \equiv \{(a, b, i, m, \tau) : m - i \leq d_n, 1 \leq m \leq n, |\tau| \leq M\}$$

and $E_{0m}(\cdot) \equiv E_0(\cdot | \mathbf{Y}_{1,m})$, $P_{\tau m}(\cdot) \equiv P_\tau(\cdot | \mathbf{Y}_{1,m})$, etc.

Lemma 3.9 *Suppose A1, A3, and A4 hold and*

$$d_n = o(n^{1/2} / \log n). \quad (3.9)$$

Then,

$$P_0 \left[\inf_{S_n} E_{0m} \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{i,m}) | X_i = a, X_{i+1} = b \right\} \geq \frac{1}{2} \right] = 1 - o(n^{-1}). \quad (3.10)$$

Proof of lemma 3.9: From (1.2), if $|\tau| \leq M$,

$$\frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{i,m}) \geq \left(\inf_{c,d} \frac{\alpha_\tau}{\alpha_0}(c, d) \right)^{m-i+1} \inf_c \frac{\pi_\tau}{\pi_0}(c) \exp \left\{ -M\delta_n \sum_{j=i}^m q_0(Y_j, M\delta_n) \right\}. \quad (a)$$

By A1 and A2, if $|\tau| \leq M$ then the first two terms are larger than $(1 - r\tau)^{m-i+1}$ for a fixed $r = r(M) < \infty$ so that

$$\begin{aligned} \inf_{S_n} E_{0m} \left\{ \frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{j,m}) | X_i = a, X_{i+1} = b \right\} & \quad (b) \\ \geq (1 + o(1)) \exp \{ -(d_n + 1)M\delta_n \max_{1 \leq j \leq n} q_0(Y_j, M\delta_n) \}. & \end{aligned}$$

But by (3.9) and A3, for some $\lambda > 0$,

$$\begin{aligned} P_0 \left[\max_{1 \leq j \leq n} q_0(Y_j, M\delta_n) \geq \log 2 / M d_n \delta_n \right] & \quad (c) \quad \text{R22: Changed} \\ \leq n P_0 [q_0(Y_1, M\delta_n) \geq \log 2 / M d_n \delta_n] & \quad \text{R23: Changed} \\ \leq n \exp \{ -\lambda (\log 2 / M) c_n \log n \} E_0 e^{\lambda q_0(Y_1, M\delta_n)}, & \end{aligned}$$

where $c_n \rightarrow \infty$ and (3.10) follows. \square

Lemma 3.10 Suppose A1 - A4 hold and $\epsilon > \frac{2}{r}$. Suppose $d_n \rightarrow \infty$, $d_n = o(n^{1/2} / (\log n)^2)$. Then

$$\sup_{S_n} \left| \frac{\alpha_{\tau,i,m}}{\alpha_{0,i,m}}(a, b) - 1 \right| = O_{1/n} \left(n^{-1/2+\epsilon} \right) \quad (3.11)$$

Proof of lemma 3.10: By lemmas 3.8 and 3.9 it is enough to show that,

$$\begin{aligned} \sup_{S_n} \{ & | E_{0m} \left(\frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{i,m}) | X_i = a, X_{i+1} = b \right) & \quad (a) \\ & - E_{0m} \left(\frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{i,m}) | X_i = a, X_{i+1} = c \right) | \} \\ = & O_{n^{-1}} \left(n^{-1/2+\epsilon} \right). \end{aligned}$$

Consider the following three Markov chains, $X'_{i+1}, \dots, X'_m; X''_{i+1}, \dots, X''_m; X'''_{i+1}, \dots, X'''_m$ where,

- i) The $\{X'_j\}$ and $\{X''_j\}$ are independent. Both have transition probabilities $\alpha_{0,j,m}$, for going from j to $j+1$, $i \leq j \leq m$, with $\mathbf{Y}_{1,m}$ held fixed, $X'_i = X''_{i+1} = a$ and $X'_{i+1} = b$, $X''_{i+1} = c$. and
- ii) $X'''_\ell = X''_\ell 1(\ell \leq T) + X'_\ell 1(\ell > T)$ where $T = \min\{\ell : X'_\ell = X''_\ell, i < \ell \leq m\} \wedge m$. Note that,

$$\{X''_\ell : i \leq \ell \leq T\} \text{ and } \{X'''_\ell : i \leq \ell \leq T\} \quad (\text{b})$$

have the same distribution. Further, if E_{0m}, P_{0m} now refer to probabilities on the space on which the data and the X'_j, X''_j, X'''_j are defined,

$$\left| E_{0m} \left(\frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{i,m}) | X_i = a, X_{i+1} = b \right) \right. \quad (\text{c})$$

$$\left. - E_{0m} \left(\frac{f_\tau}{f_0}(\mathbf{X}_{i,m}, \mathbf{Y}_{i,m}) | X_i = a, X_{i+1} = c \right) \right|$$

$$= \left| E_{0m} \left(\frac{f_\tau}{f_0}(\mathbf{X}'_{i,m}, \mathbf{Y}_{i,m}) - \frac{f_\tau}{f_0}(\mathbf{X}''_{i,m}, \mathbf{Y}_{i,m}) \right) \right|$$

$$= \left| E_{0m} \left[\left(\frac{f_\tau}{f_0}(\mathbf{X}'_{i,T}, \mathbf{Y}_{i,T}) - \frac{f_\tau}{f_0}(\mathbf{X}'''_{i,T}, \mathbf{Y}_{i,T}) \right) \right. \right. \quad (\text{d})$$

$$\left. \cdot \frac{\pi_0}{\pi_\tau}(X'_{T+1}) \frac{f_\tau}{f_0}(X'_{T+1}, \dots, X'_m, Y_{T+1}, \dots, Y_m) \frac{\alpha_\tau}{\alpha_0}(X'_T, X'_{T+1}) \right] \right|.$$

By A1 and A2, for $|\tau| \leq M$, d_n as above there exists $c(M) < \infty$ such that, if $A_n \equiv \max\{q_0(Y_j, M\delta_n) : 1 \leq j \leq n\}$,

$$\exp\{-\delta_n(T-i)(MA_n + c)\} \quad (\text{e})$$

$$\leq \frac{f_\tau}{f_0}(\mathbf{X}'_{i,T}, \mathbf{Y}_{i,T})$$

$$\leq \exp\{\delta_n(T-i)(MA_n + c)\}$$

The same holds if $\mathbf{X}'_{i,T}$ is replaced by $\mathbf{X}'''_{i,T}$ and also

$$\frac{f_\tau}{f_0}(\mathbf{X}'_{T+1,m}, \mathbf{Y}_{T+1,m}) \leq \exp\{\delta_n d_n(MA_n + c)\}. \quad (\text{f})$$

By A3 and (c) of the proof of lemma 3.9

$$A_n = O_{n^{-1}} \left((\log n)^2 \right). \quad (\text{g})$$

Then, from (d), (e), (f), and (g), (a) follows if

$$\sup_{S_n} \{E_{0m}(e^{(T-i)a_n} - e^{-(T-i)a_n})\} = O_{n-1} \left(n^{-1/2+\epsilon} \right) \quad (\text{h})$$

for

$$a_n = O(\delta_n (\log n)^2). \quad (\text{i})$$

Now,

$$P_{0m}[T > i + t] \leq \prod_{j=i+1}^{i+t} (1 - K\mu_0^2(Y_j)). \quad (\text{j})$$

since for $j \geq i$

$$\begin{aligned} P_{0m}[X'_{j+1} = X''_{j+1} | X'_j = a, X''_j = b] &= \sum_c \alpha_{0,j,m}(a, c) \alpha_{0,i,m}(b, c) \quad (\text{k}) \\ &\geq K\mu_0^2(Y_{j+1}) \end{aligned}$$

by lemma 3.1. But, by A4

$$\begin{aligned} P_0 \left[\min_{1 \leq j \leq n} \{K\mu_0^2(Y_j)\} \leq b_n \right] &= P_0 \left[\max_{1 \leq j \leq n} \{\rho_0(Y_j)\} \geq \frac{\gamma^2((K/b_n)^{1/2} - 1)}{K - 1} \right] \quad (\text{l}) \\ &= o(n^{-1}) \end{aligned}$$

if

$$b_n = o(n^{-2/r}). \quad (\text{m})$$

Note that for any integer valued random variable $N \geq 1$

$$Ea^N = a + \sum_{t=1}^{\infty} (a^{t+1} - a^t) P[N > t] \quad (\text{n})$$

From (j), (l), (n), if $b_n = o(n^{-2/r})$, $b_n n^\epsilon / (\log n)^2 \rightarrow \infty$, then $a_n = o(b_n)$ and, R24: Changed with probability $1 - o(n^{-1})$,

$$\begin{aligned} &\max\{E_{0m}(e^{(T-i)a_n} - e^{-(T-i)a_n}) : m - i \leq d_n, 1 \leq m \leq n\} \\ &\leq e^{a_n} - e^{-a_n} + \sum_{t=1}^{\infty} (e^{a_n} - 1) e^{t(a_n - b_n)} \\ &= e^{a_n} - e^{-a_n} + (e^{a_n} - 1) e^{a_n - b_n} (1 - e^{(a_n - b_n)})^{-1} \\ &= O(a_n (b_n - a_n)^{-1}) \\ &= O(a_n b_n^{-1}) \end{aligned}$$

and (a) follows from (h). □

Lemma 3.11 *If A1 - A4 hold, $\epsilon > \frac{2}{r}$, then*

$$\sup_{S_n} |P_{\tau m}[X_m = a] - P_{0m}[X_m = a]| = O_{1/n} \left(n^{-1/2+2\epsilon} \right) \quad (3.12)$$

Proof of lemma 3.11: For fixed a let $V_{\tau, \ell, m} \in R^K$ be the column vector with coordinates:

$$V_{\tau, \ell, m}(\cdot) = P_{\tau m}[X_m = a | X_\ell = \cdot], \quad \ell \leq m. \quad (a)$$

Then,

$$V_{\tau, \ell, m} = \alpha_{\tau, \ell, m} \dots \alpha_{\tau, m-1, m} V_{\tau, m, m}. \quad (b)$$

By lemma 3.3

$$\begin{aligned} & \sup \{ |V_{\tau, \ell, m}(b) - V_{\tau, \ell, m}(c)| : b, c, |\tau| < M \} \\ & \leq \prod_{j=\ell+1}^{m-1} (1 - 2\mu_0(Y_j)) \\ & \leq e^{-(m-\ell-1)B_n} \end{aligned} \quad (c)$$

where,

$$B_n = 2 \min_{1 \leq j \leq n} \mu_0(Y_j). \quad (d)$$

Then,

$$\sup \{ |V_{\tau, \ell, m}(b) - V_{\tau, \ell, m}(c)| e^{(m-\ell-1)b_n} : b, c, |\tau| \leq M, \ell \leq m \} = O_{n^{-1}}(1) \quad (e)$$

if $b_n = o(n^{-\frac{2}{r}})$, by arguing as in (l) of lemma 3.10. Therefore, if $c_{\tau, \ell, m} = K^{-1} \sum_b V_{\tau, \ell, m}(b)$ then

$$\sup \{ \|V_{\tau, \ell, m} - c_{\tau, \ell, m} \mathbf{1}\| e^{b_n(m-\ell-1)} : m, \ell, |\tau| \leq M \} = O_{n^{-1}}(1) \quad (f)$$

where $\|\cdot\|$ is the L_∞ on R^k and $\mathbf{1}$ is the vector of 1's. Then from (b),

$$\begin{aligned} \|V_{\tau, \ell, m} - V_{0, \ell, m}\| &= \|\alpha_{\tau, \ell, m} V_{\tau, \ell+1, m} - \alpha_{0, \ell, m} V_{0, \ell+1, m}\| \\ &\leq \|(\alpha_{\tau, \ell, m} - \alpha_{0, \ell, m}) V_{\tau, \ell+1, m}\| + \|V_{\tau, \ell+1, m} - V_{0, \ell+1, m}\|. \end{aligned} \quad (g)$$

Further, from lemma 3.10, if $m - \ell = o(n^{1/2}/(\log n)^2)$, $b_n = o(n^{-2/r})$

$$\|(\alpha_{\tau,\ell,m} - \alpha_{0,\ell,m})V_{\tau,\ell+1,m}\| \leq \rho_n e^{-(m-\ell-1)b_n} \quad (\text{h})$$

where $\rho_n = O_{n^{-1}}(c_n)$, $c_n = n^{-1/2+\epsilon}$ since

$$(\alpha_{\tau,\ell,m} - \alpha_{0,\ell,m})\mathbf{1} = 0. \quad (\text{i})$$

Iterating (g) and using (h), we get, if $d_n = o(n^{1/2}/(\log n)^2)$, $b_n = o(n^{-2/r})$

$$\sup\{\|V_{\tau,\ell,m} - V_{0,\ell,m}\| : m - \ell \leq d_n, |\tau| \leq M\} = O_{n^{-1}}(c_n b_n^{-1}). \quad (\text{j})$$

Finally,

$$\begin{aligned} & |P_{\tau m}[X_m = a] - P_{0m}[X_m = a]| \quad (\text{k}) \\ &= \left| \sum_b \{P_{\tau m}[X_\ell = b]V_{\tau,\ell,m}(b) - P_{0m}[X_\ell = b]V_{0,\ell,m}(b)\} \right| \\ &\leq \left| \sum_b (P_{\tau m}[X_\ell = b] - P_{0m}[X_\ell = b])V_{\tau,\ell,m}(b) \right| + \|V_{\tau,\ell,m} - V_{0,\ell,m}\|. \end{aligned}$$

By (f) the first term in (k) is, if $m - \ell \geq d_n$, $= O_{n^{-1}}(e^{-d_n b_n})$. If we use (j) and put $b_n = n^{-\epsilon}$, $d_n = n^\epsilon(\log n)^2$ the lemma follows. \square

Lemma 3.12 *Under A1 - A4, if $k = o(n^{1/2-\gamma})$, for some $\gamma > 0$,*

$$\begin{aligned} & \sup \left\{ \left| \frac{\ell_\tau(\mathbf{Y}_m^{(k)} | X_{mk+1})}{\ell_0} - 1 \right| : |\tau| \leq M, 1 \leq m \leq N \right\} \quad (3.13) \\ &= O_{1/n}(n^{-\gamma/2}) \end{aligned}$$

Proof of lemma 3.12: Note that for any $p > 1$,

$$\begin{aligned} & E_0 \sup \left\{ \left| \frac{\ell_\tau(\mathbf{Y}_0^{(k)} | X_1)}{\ell_0} - 1 \right|^p : |\tau| \leq M \right\} \quad (\text{a}) \\ &= E_0 \sup \left\{ \left| E_0 \left[\left(\frac{\pi_0}{\pi_\tau}(X_1) \frac{f_\tau(\mathbf{X}_0^{(k)}, \mathbf{Y}_0^{(k)})}{f_0} - 1 \right) \middle| X_1, \mathbf{Y}_0^{(k)} \right] \right|^p : |\tau| \leq M \right\} \\ &\leq (1 + o(1)) E_0 \sup \left\{ \left| \frac{f_\tau(\mathbf{X}_0^{(k)}, \mathbf{Y}_0^{(k)})}{f_0} - 1 \right|^p : |\tau| \leq M \right\} + o(1) \end{aligned}$$

But, for any differentiable function $A(\vartheta)$ with $A(0) = 0$,

$$\begin{aligned} \sup \left\{ \left| e^{A(\vartheta)} - 1 \right|, |\vartheta| \leq \delta \right\} &\leq \delta \sup \left\{ |A'(\vartheta)| e^{A(\vartheta)} : |\vartheta| \leq \delta \right\} \\ &\leq \delta M_\delta e^{\delta M_\delta} \end{aligned} \quad (\text{b})$$

where $M_\delta \equiv \sup \{|A'(\vartheta)| : |\vartheta| \leq \delta\}$. We conclude that,

$$\begin{aligned} E_0 \sup \left\{ \left| \frac{f_\tau}{f_0}(\mathbf{X}_0^{(k)}, \mathbf{Y}_0^{(k)}) - 1 \right|^p : |\tau| \leq M \right\} \\ \leq (M\delta_n)^p E_0 \left\{ \left(\sum_{j=1}^k \tilde{q}(Y_j, M\delta_n) \right)^p \exp \left[pM\delta_n \sum_{j=1}^k \tilde{q}(Y_j, M\delta_n) \right] \right\}, \end{aligned} \quad (\text{c})$$

where

$$\tilde{q}(y, \delta) = q(y, \delta) + \sup \{ |\nabla \log \alpha_\vartheta(a, b)| : |\vartheta - \vartheta_0| < \delta, a, b \}. \quad (\text{d})$$

Bound the right hand side of (c) by,

$$(M\delta_n)^p E_0^{\frac{1}{1+\epsilon}} \left\{ \left(\sum_{j=1}^k \tilde{q}(Y_j, M\delta_n) \right)^{p(1+\epsilon)} \right\} E_0^{\frac{\epsilon}{1+\epsilon}} \left\{ \exp \left[\frac{p(1+\epsilon)M\delta_n}{\epsilon} \sum_{j=1}^k \tilde{q}(Y_j, M\delta_n) \right] \right\} \quad (\text{e})$$

The second term in (e) is bounded by

$$k^p \left[E_0 \left\{ \tilde{q}^{p(1+\epsilon)}(Y_1, M\delta_n) \right\} \right]^{\frac{1}{1+\epsilon}} \quad (\text{f})$$

and use A3 to bound the third by,

$$\begin{aligned} &\left[\max_a E_0 \left\{ \exp \left(\frac{p(1+\epsilon)}{\epsilon} M\delta_n q(Y_1, M\delta_n) \right) \middle| X_1 = a \right\} \right]^{k\epsilon/(1+\epsilon)} \\ &= \left(1 + \frac{O(1)}{\sqrt{n}} \right)^{k\epsilon/(1+\epsilon)} \\ &= 1 + o(1) \end{aligned} \quad (\text{g})$$

since $k = o(n^{1/2})$ and $\delta_n \rightarrow 0$. Therefore,

$$\begin{aligned} P_0 \left[\sup_{\substack{1 \leq m \leq N \\ |\tau| \leq M}} \left\{ \left| \frac{\ell_\tau}{\ell_0}(\mathbf{Y}_m^{(k)} | X_{mk+1}) - 1 \right| \right\} \geq n^{-\gamma/2} \right] \\ \leq O(1) \frac{n}{k} (k\delta_n)^p n^{p\gamma/2} \\ = o(n^{-1}) \end{aligned} \quad (\text{h})$$

if $k = O(n^{1/2-\gamma})$, $p > 2 + 3/\gamma$.

□

Lemma 3.13 *Under A1 - A4 if $k = o(n^{1/2-\gamma})$, for some $\gamma > 0$ and $\epsilon > \frac{2}{r}$*

$$\sup \left\{ \frac{|L_{\tau m} - L_{\tau m}^*|}{L_{\tau m}^*} : 1 \leq m \leq N, |\tau| \leq M \right\} = O_{1/n} \left(n^{-\frac{1}{2}+2\epsilon} \right). \quad (3.14)$$

Proof of lemma 3.13: By (3.1)

$$\min_a \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)B_\tau(\mathbf{Y}_{1,mk})}{\ell_0} \leq L_{\tau m} \leq \max_a \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)B_\tau(\mathbf{Y}_{1,mk})}{\ell_0} \quad (a)$$

where,

$$B_\tau(\mathbf{Y}_{1,mk}) = \frac{\sum_a P_\tau[X_{mk+1} = a | \mathbf{Y}_{1,mk}] \ell_0(\mathbf{Y}_m^{(k)}|a)}{\sum_a P_0[X_{mk+1} = a | \mathbf{Y}_{1,mk}] \ell_0(\mathbf{Y}_m^{(k)}|a)}. \quad (b)$$

But,

$$|B_\tau(\mathbf{Y}_{1,mk}) - 1| \leq \max_a \left| \frac{P_\tau[X_{mk+1} = a | \mathbf{Y}_{1,mk}]}{P_0[X_{mk+1} = a | \mathbf{Y}_{1,mk}]} - 1 \right|. \quad (c)$$

It follows from lemmas 3.11 and 3.4,

$$\sup \{ |B_\tau(\mathbf{Y}_{1,mk}) - 1| : |\tau| \leq M, 1 \leq m \leq N \} = O_{n^{-1}} \left(n^{-1/2+2\epsilon} \right). \quad (d)$$

On the other hand

$$\frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)}{\ell_0} = \frac{\pi_0}{\pi_\tau} E_0 \left\{ \frac{f_\tau(\mathbf{X}_m^{(k)}, \mathbf{Y}_m^{(k)})}{f_0} \middle| X_{mk+1} = a, \mathbf{Y}_m^{(k)} \right\} \quad (e)$$

so that by (a) of the proof of lemma 3.10, if $k = o(n^{1/2}/(\log n)^2)$

$$\sup \left\{ \left| \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)}{\ell_0} - \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|b)}{\ell_0} \right| : m, |\tau| \leq M, a, b \right\} = O_{n^{-1}} \left(n^{-1/2+\epsilon} \right). \quad (f)$$

From (a), (d), (f), and lemma 3.12 we obtain lemma 3.13.

□

Lemma 3.14 *Under A1 - A5*

$$E_0 \sum_{m=1}^N |L_{\tau m}^* - 1| = O\left(\left(\frac{n}{k}\right)^{1/2}\right) \quad (3.15)$$

Proof of lemma 3.14: Note that

$$E_0\{|L_{\tau 0}^* - 1||X_1 = a\} = \|\mathcal{L}_\tau(\mathbf{Y}_{1,k}|X_1 = a) - \mathcal{L}_0(\mathbf{Y}_{1,k}|X_1 = a)\| \quad (a)$$

where $\|\cdot\|$ denotes variational distance. Therefore

$$E_0\{|L_{\tau 0}^* - 1||X_1 = a\} \quad (b)$$

$$\leq \|\mathcal{L}_\tau((\mathbf{X}_{1,k}, \mathbf{Y}_{1,k})|X_1 = a) - \mathcal{L}_0((\mathbf{X}_{1,k}, \mathbf{Y}_{1,k})|X_1 = a)\| \quad (c)$$

$$\leq 2H(\mathcal{L}_{0a}, \mathcal{L}_{1a}) \left(2 - H^2(\mathcal{L}_{0a}, \mathcal{L}_{1a})\right)^{1/2}$$

where $\mathcal{L}_{0a}, \mathcal{L}_{1a}$ are the laws in (c) and H is Hellinger distance, by a standard inequality (LeCam (1986) p. 47). But,

$$1 - H^2(\mathcal{L}_{0a}, \mathcal{L}_{1a}) \quad (d)$$

$$= E_0 \left\{ \left(\frac{f_\tau}{f_0} \right)^{\frac{1}{2}} (\mathbf{X}_{1,k}, \mathbf{Y}_{1,k}) | X_1 = a \right\}$$

$$= E_0 \left\{ \left(\frac{\pi_\tau}{\pi_0} \right)^{1/2} (a) \prod_1^{k-1} \left(\frac{\alpha_\tau}{\alpha_0} \right)^{1/2} (X_i, X_{i+1}) \right.$$

$$\left. \prod_{i=1}^k E_0 \left[\left(\frac{g_\tau}{g_0} \right)^{1/2} (Y_i | X_i) \right] \middle| X_1 = a \right\}.$$

But

$$\prod_{i=1}^k E_0 \left[\left(\frac{g_\tau}{g_0} \right)^{1/2} (Y_i | X_i) \right] \quad (e)$$

$$= \prod_{i=1}^k E_0 \left\{ e^{\frac{1}{2} \log(g_\tau/g_0)(Y_i | X_i)} \right\}$$

$$\geq \prod_{i=1}^k \left[1 - \frac{\delta_n^2}{2} |\tau|^2 p E_0 \left\{ \left(\frac{1}{4} q_0^2(Y_i, M\delta_n) + \frac{1}{2} q_{02}(Y_i, M\delta_n) \right) e^{\frac{|\tau|}{2} \delta_n q_0(Y_i, M\delta_n)} \right\} \right]$$

$$\geq 1 - O\left(\frac{k}{n}\right)$$

by Taylor expansion and A3 and A5. Similarly, by A1 and A2:

$$E_0 \left\{ \left(\frac{\pi_\tau}{\pi_0} \right)^{1/2} (a) \prod_1^{k-1} \left(\frac{\alpha_\tau}{\alpha_0} \right)^{1/2} (X_i, X_{i+1}) \middle| X_1 = a \right\} \geq 1 - O\left(\frac{k}{n}\right). \quad (\text{f})$$

Finally, we conclude from (e) and (f):

$$\sum_{a=1}^K H^2(\mathcal{L}_{0a}, \mathcal{L}_{1a}) \pi_0(a) = O\left(\frac{k}{n}\right). \quad (\text{g})$$

The lemma is proved by (a), (b), and (g). \square

Lemma 3.15 *If $L_{\tau m}^{(d)}$ is given by (2.2), $r > 8$, $k = o(n^{1/2-\gamma})$ for some $\gamma > 0$, $d = n^\epsilon \log^2 n$, $\epsilon > \frac{2}{r}$ then*

$$\sup \left\{ \left| L_{\tau m}^{(d)} - L_{\tau m} \right| : |\tau| \leq M, 1 \leq m \leq N \right\} = O_{n^{-1}} \left(n^{-1-\epsilon} \right). \quad (3.16)$$

Proof of lemma 3.15: By lemma 3.2, if B_n is given by (d) of lemma 3.11,

$$\begin{aligned} & \sup_{\substack{1 \leq m \leq N \\ |\tau| \leq M}} \left\{ \left| P_\tau[X_{mk+1} = a | \mathbf{Y}_{mk-d, mk}] - P_\tau[X_{mk+1} = a | \mathbf{Y}_{1, mk}] \right| \right\} \quad (\text{a}) \\ & \leq \max_{m \leq 1} \left\{ \sum_{\ell=1}^{mk-d-1} \prod_{j=\ell+1}^{mk} (1 - 2\mu_0(Y_j)) \right\} \\ & \leq \frac{e^{-(d-1)B_n}}{1 - e^{-B_n}} \\ & = O_{1/n} \left(n^{-1-2\epsilon} \right) \end{aligned}$$

by arguing as for (l) of lemma 3.10. But, by lemma 3.1,

$$\begin{aligned} P_0[X_\ell = a | \mathbf{Y}_{1, n}] & \geq \min_{a, b} \alpha_{0, \ell-1, n}(a, b) \quad (\text{b}) \\ & \geq \min\{\mu_0(Y_j) : 1 \leq j \leq n\} \end{aligned}$$

and, hence

$$P_0[\min_{a, \ell} P_0[X_\ell = a | \mathbf{Y}_{1, n}] \geq n^{-\epsilon}] = 1 - o(n^{-1}). \quad (\text{c})$$

But, arguing as for lemma 3.13,

$$|L_{\tau m} - L_{\tau m}^{(d)}| \leq A_m(\tau) \max_a \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)}{\ell_0} + A_m(0)L_{\tau m} \quad (d)$$

where

$$A_m(\tau) \equiv \max_a \left\{ \frac{|P_\tau[X_{mk+1} = a | \mathbf{Y}_{1,mk}] - P_\tau[X_{mk+1} = a | \mathbf{Y}_{mk-d,mk}]|}{P_0[X_{mk+1} = a | \mathbf{Y}_{mk-d,mk}]} \right\}. \quad (e)$$

By (a) and (c),

$$\sup\{A_m(\tau) : m, |\tau| \leq M\} = O_{n^{-1}}(n^{-(1+\epsilon)}) \quad (f)$$

and lemma 3.15 follows from (d), lemma 3.12, and lemma 3.13. \square

Lemma 3.16 *Suppose A1 - A4 hold. Let $d = n^\epsilon \log^2 n$, $\epsilon > \frac{2}{r}$, and $k = n^{4\epsilon+\gamma}$ for some $\gamma > 0$, $4\epsilon + \gamma < \frac{1}{2}$, so that $r > 16$. Then*

$$E_0 \left\{ \left(\sum_m (L_{\tau m}^{(d)} - L_{\tau m}^*) \right)^2 \right\} = O(n^{-\gamma}) \quad (3.17)$$

Proof of lemma 3.16: For any fixed u , we first bound

$$E_0 \left\{ (L_{\tau m}^{(d)})^u \right\} \leq E_0 \left\{ \left(\max_a \frac{P_\tau}{P_0}[X_{mk+1} = a | \mathbf{Y}_{mk-d,mk}] \max_a \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)}{\ell_0} \right)^u \right\}. \quad (a)$$

Now the first term in (a) is uniformly bounded by lemma 3.4. The second is bounded by,

$$\exp\left\{ M \delta_n u \sum_{j=mk+1}^{(m+1)k} q_0(Y_j, M \delta_n) \right\}. \quad (b)$$

Thus, if $k = o(n^{1/2})$, by A3, for all u , eventually,

$$E_0(L_{\tau m}^{(d)})^u \leq C_1(1 + C_2 \delta_n)^k \leq C_3. \quad (c)$$

Similarly,

$$E_0(L_{\tau m}^*)^u \leq C_4. \quad (\text{d})$$

Now,

$$\begin{aligned} & |L_{\tau m}^{(d)} - L_{\tau m}^*| \quad (\text{e}) \\ & \leq \max_{a,b} \left| \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)}{\ell_0} - \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|b)}{\ell_0} \right| \\ & \quad + \max_a \frac{\ell_\tau(\mathbf{Y}_m^{(k)}|a)}{\ell_0} \max_a \left| \frac{P_\tau}{P_0}[X_{mk+1} = a | \mathbf{Y}_{mk-d, mk}] - 1 \right| \\ & = O_{n-1}(n^{-1/2+\epsilon}) + O_{n-1}(n^{-1/2+2\epsilon}) \end{aligned}$$

by (f) of lemma 3.13, lemma 3.12, lemma 3.4 and lemma 3.11 and (a) of lemma 3.15. Let $c_n = cn^{-1/2+2\epsilon}$ for some large enough c . Note that

$$\begin{aligned} & E_0 |L_{\tau m}^{(d)} - L_{\tau m}^*|^{2+8\epsilon} \quad (\text{f}) \\ & \leq c_n^{2+8\epsilon} + E_0 \left\{ |L_{\tau m}^{(d)} - L_{\tau m}^*|^{2+8\epsilon} 1(|L_{\tau m}^{(d)} - L_{\tau m}^*| \geq c_n) \right\} \\ & \leq c_n^{2+8\epsilon} + E_0^{16\epsilon^2} \left\{ |L_{\tau m}^{(d)} - L_{\tau m}^*|^{(2+8\epsilon)/16\epsilon^2} \right\} P_0^{1-16\epsilon^2} \left\{ |L_{\tau m}^{(d)} - L_{\tau m}^*| > c_n \right\} \\ & \leq 2c_n^{2+8\epsilon} \end{aligned}$$

for large enough n .

We will apply the Ibragimov-Linnik lemma 3.7, with $\delta = 8\epsilon$. Note that, if $d \leq k$, by the geometric ergodicity of the chain under A1, A2, the variational norm distance between the joint distribution of $(L_{\tau m_1}^{(d)} - L_{\tau m_1}^*, L_{\tau m_2}^{(d)} - L_{\tau m_2}^*)$ and the product of the marginals is bounded by $C\xi^{|m_1 - m_2|}$ for some $C < \infty$, $\xi < 1$ and all m_1, m_2 . Hence, using (f) above

$$\begin{aligned} E_0 \left\{ \left(\sum_{m=1}^N (L_{\tau m}^{(d)} - L_{\tau m}^*) \right)^2 \right\} & = O\left(\frac{n}{k} c_n^2\right) \quad (\text{g}) \\ & = O(n^{-\gamma}) \end{aligned}$$

under our conditions on k, c_n .

□

Proof of lemma 2.1: It's enough to show all terms on the RHS of (2.6) are $O_{e_n}(n^{-\gamma/2}/e_n)$. The first term is equal to

$$\sum_{m=1}^N (L_{\tau m} - L_{\tau m}^{(d)}) + \sum_{m=1}^N (L_{\tau m}^{(d)} - L_{\tau m}^*) = O_{e_n}(n^{-\gamma/2}e_n^{-1/2}) \quad (\text{a})$$

by lemma 3.15 and lemma 3.16. The second term can be bounded by

$$\begin{aligned} \sup_{\substack{1 \leq m \leq N \\ |\tau| \leq M}} \left\{ \frac{|L_{\tau m} - L_{\tau m}^*|}{L_{\tau m}^*} \right\} \sum_{m=1}^N |L_{\tau m}^* - 1| &= O_{e_n} \left(n^{-1/2+2\epsilon} \left(\frac{n}{k} \right)^{1/2} / e_n \right) \quad (\text{b}) \\ &= O_{e_n}(n^{-\gamma/2}/e_n) \end{aligned}$$

by lemmas 3.13 and 3.14. Finally the third term is negligible since

$$\begin{aligned} |R_n| &\leq \left(1 - \sup \left\{ \frac{|L_{\tau m} - L_{\tau m}^*|}{L_{\tau m}^*} : 1 \leq m \leq N, |\tau| \leq M \right\} \right)^{-2} \quad (\text{c}) \\ &= O_{n^{-1}}(1) \end{aligned}$$

and

$$\sum_{m=1}^N \left(\frac{|L_{\tau m} - L_{\tau m}^*|}{L_{\tau m}^*} \right)^2 = O_{1/n}(n^{-\gamma}) \quad (\text{d})$$

both by lemma 3.13. □

Proof of lemma 2.2: Expand

$$\begin{aligned} \log L_{\tau m}^* &= \delta_n \tau^T \nabla \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \quad (\text{a}) \\ &+ \frac{1}{2n} \tau^T \left\| \frac{\partial^2}{\partial \vartheta_i \partial \vartheta_j} \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \right\| \tau \\ &+ \delta_n^3 \int_0^1 \frac{(1-\lambda)^2}{2} \sum_{a,b,c} \tau_a \tau_b \tau_c \frac{\partial^3}{\partial \vartheta_a \partial \vartheta_b \partial \vartheta_c} \log \ell_{\lambda \tau}(\mathbf{Y}_m^{(k)} | X_{mk+1}) d\lambda. \end{aligned}$$

We use a classical formula based on lemma 3.6. If \mathbf{B} is generated by X_{mk+1} , $\mathbf{Y}_m^{(k)}$, and we suppress arguments in f_ϑ ,

$$\begin{aligned}
& \frac{\partial^3}{\partial\vartheta_a\partial\vartheta_b\partial\vartheta_c} \log \ell_\vartheta(\mathbf{Y}_m^{(k)}|X_{mk+1}) \tag{b} \\
&= E_\vartheta \left\{ \frac{\partial^3}{\partial\vartheta_a\partial\vartheta_b\partial\vartheta_c} \log f_\vartheta | \mathbf{B} \right\} + \text{cov}_\vartheta \left\{ \frac{\partial^2}{\partial\vartheta_a\partial\vartheta_b} \log f_\vartheta, \frac{\partial}{\partial\vartheta_c} \log f_\vartheta | \mathbf{B} \right\} \\
&+ \text{cov}_\vartheta \left\{ \frac{\partial^2}{\partial\vartheta_a\partial\vartheta_c} \log f_\vartheta, \frac{\partial}{\partial\vartheta_b} \log f_\vartheta | \mathbf{B} \right\} + \text{cov}_\vartheta \left\{ \frac{\partial^2}{\partial\vartheta_b\partial\vartheta_c} \log f_\vartheta, \frac{\partial}{\partial\vartheta_a} \log f_\vartheta | \mathbf{B} \right\} \\
&- \text{cov}_\vartheta \left\{ \frac{\partial}{\partial\vartheta_a} \log f_\vartheta, \frac{\partial}{\partial\vartheta_b} \log f_\vartheta, \frac{\partial}{\partial\vartheta_c} \log f_\vartheta | \mathbf{B} \right\} \\
&- \text{cov}_\vartheta \left\{ \frac{\partial}{\partial\vartheta_a} \log f_\vartheta, \frac{\partial}{\partial\vartheta_c} \log f_\vartheta, \frac{\partial}{\partial\vartheta_b} \log f_\vartheta | \mathbf{B} \right\} \\
&- \text{cov}_\vartheta \left\{ \frac{\partial}{\partial\vartheta_b} \log f_\vartheta, \frac{\partial}{\partial\vartheta_c} \log f_\vartheta, \frac{\partial}{\partial\vartheta_a} \log f_\vartheta | \mathbf{B} \right\} - \frac{\partial^3}{\partial\vartheta_a\partial\vartheta_b\partial\vartheta_c} \log \pi_\vartheta(X_{mk+1}).
\end{aligned}$$

We see from (b) and assumptions A1 and A2 that to bound the third term in (a) it suffices to bound, for $|\vartheta - \vartheta_0| \leq M\delta_n$, all a, b, c

$$E_0 \left\{ E_\vartheta \left[\left| \sum_{j=1}^k \frac{\partial^3}{\partial\vartheta_a\partial\vartheta_b\partial\vartheta_c} \log g_\vartheta(Y_j|X_j) \right| | Y_1, \dots, Y_k \right] \right\}, \tag{c}$$

$$\begin{aligned}
& E_0 \left\{ E_\vartheta \left[\left| \sum_{j=1}^k \frac{\partial^2}{\partial\vartheta_a\partial\vartheta_b} \log g_\vartheta(Y_j|X_j) \right| \right. \right. \\
& \quad \left. \left. \cdot \left(1 + \left| \sum_{j=1}^k \frac{\partial}{\partial\vartheta_c} \log g_\vartheta(Y_j|X_j) \right| \right) | Y_1, \dots, Y_k \right] \right\}, \tag{d}
\end{aligned}$$

and

$$E_0 \left\{ E_\vartheta \left[\left| \sum_{j=1}^k \frac{\partial}{\partial\vartheta_a} \log g_\vartheta(Y_j|X_j) \right|^3 | Y_1, \dots, Y_k \right] \right\}. \tag{e}$$

We can apply lemma 3.5 to all of these and use A3 to conclude that, under A5, (c)–(e) are uniformly $O(k^3)$. To do so we take r in the lemma as close to

1 as possible and s and t as large as necessary since, by A3, and by arguing as in lemma 3.16 (b), $E_0 \exp |t\Lambda| < \infty$ for all $k = o(n^{1/2})$, t . Therefore, the expectation of the remainder in (a) is $O(n^{-3/2}k^3)$. The lemma follows since there are n/k terms like that in the LHS of (2.10). under our assumptions. \square

R25: Changed
My stupid mistake of the last revision was forgetting the need for a sentence like the last one.

Lemma 3.17 *Let $-k \geq -j + 2$ and*

$$S(j, k) \equiv \max_{a,b,c} \left\{ \left| P_0[X_{-k} = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = b] - P_0[X_{-k} = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = c] \right| \right\} \quad (3.18)$$

Then,

$$S(j, k) \leq 2\gamma^{-1}(\vartheta_0) \prod_{i=-k+1}^0 (1 - 2\mu_0(Y_i)) \quad (3.19)$$

Proof of lemma 3.17:

$$\begin{aligned} & P_0[X_{-k} = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = b] \quad (a) \\ &= \frac{P_0[X_1 = b | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_{-k} = a]}{P_0[X_1 = b | X_{-j+2}, \mathbf{Y}_{-j+2,0}]} P_0[X_{-k} = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}]. \end{aligned}$$

Then

$$S(j, k) \leq 2 \max_{a,b} \left\{ \left| \frac{P_0[X_1 = b | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_{-k} = a]}{P_0[X_1 = b | X_{-j+2}, \mathbf{Y}_{-j+2,0}]} - 1 \right| \right\}. \quad (b)$$

But,

$$\begin{aligned} & P_0[X_1 = b | X_{-j+2}, \mathbf{Y}_{-j+2,0}] \quad (c) \\ &= \sum_c P_0[X_1 = b | X_{-k} = c, \mathbf{Y}_{-k+1,0}] P_0[X_{-k} = c | X_{-j+2}, \mathbf{Y}_{-j+2,0}] \end{aligned}$$

and hence,

$$\begin{aligned} & S(j, k) \quad (d) \\ & \leq 2 \max_{a,b} \frac{\sum_c |P_0[X_1 = b | X_{-k} = c, \mathbf{Y}_{-k+1,0}] - P_0[X_1 = b | X_{-k} = a, \mathbf{Y}_{-k+1,0}]|}{\min_b P_0[X_1 = b | X_{-j+2}, \mathbf{Y}_{-j+2,0}]} \\ & \leq 2\gamma^{-1}(\vartheta_0) K \prod_{j=-k+1}^0 (1 - 2\mu_0(Y_j)) \end{aligned}$$

by lemma 3.3 and 3.4

□

Proof of lemma 2.3: Without loss of generality take $\vartheta_0 = 0$. Write,

$$\ell_{\vartheta}(Y_1, \dots, Y_k | X_1) = \prod_{j=1}^k \frac{g_{j\vartheta}}{g_{(j-1)\vartheta}}(X_1, \mathbf{Y}_{1,j}) \quad (\text{a})$$

where $g_{j\vartheta}(X_1, \mathbf{Y}_{1,j})$ is the joint density of $(X_1, \mathbf{Y}_{1,j})$, for $j \geq 1$, and $g_{0\vartheta} = \pi_{\vartheta}(X_1)$. Take $\dim(\vartheta) = 1$. The generalization is trivial. Then

$$\begin{aligned} & \frac{\partial}{\partial \vartheta} \log \ell_{\vartheta}(\mathbf{Y}_0^{(k)} | X_1) \\ &= \sum_{j=1}^k \left[\frac{\partial}{\partial \vartheta} \log g_{j\vartheta}(X_1, \mathbf{Y}_{1,j}) - \frac{\partial}{\partial \vartheta} \log g_{(j-1)\vartheta}(X_1, \mathbf{Y}_{1,j-1}) \right]. \end{aligned} \quad (\text{b})$$

The terms in brackets are of course martingale summands and we arrive at the identity,

R26: Changed

$$\begin{aligned} & E_0 \left\{ \left(\frac{\partial}{\partial \vartheta} \log \ell_0(\mathbf{Y}_0^{(k)} | X_1) \right)^2 \right\} \\ &= \sum_{j=1}^k E_0 \left\{ \left(\frac{\partial}{\partial \vartheta} \log g_{j0}(X_1, \mathbf{Y}_{1,j}) - \frac{\partial}{\partial \vartheta} \log g_{(j-1)0}(X_1, \mathbf{Y}_{1,j-1}) \right)^2 \right\} \\ &= \sum_{j=1}^k E_0 \{ U_j^2(X_1, \mathbf{Y}_{1,j}) \}, \quad \text{say} \\ &= \sum_{j=1}^k E_0 \{ U_j^2(X_{-j+2}, \mathbf{Y}_{-j+2,1}) \} \end{aligned} \quad (\text{c})$$

where (X_j, Y_j) , $-\infty < j < \infty$ is the two sided stationary sequence such that (X_j, Y_j) , $j \geq 1$ are distributed according to P_{ϑ} . We claim that,

$$E_0 \{ U_j^2(X_{-j+2}, \mathbf{Y}_{-j+2,1}) \} \rightarrow I(\vartheta_0) \quad (\text{d})$$

and that combined with (c) clearly establishes (2.11). Now, if we use $\frac{b'}{b}(\vartheta)$ for $\frac{\partial}{\partial \vartheta} \log b(\vartheta)$,

$$\begin{aligned}
& U_j(X_{-j+2}, \mathbf{Y}_{-j+2,1}) \tag{e} \\
&= E_0 \left\{ \sum_{m=-j+2}^1 \frac{g'_0(Y_m|X_m)}{g_0} + \sum_{m=-j+2}^0 \frac{\alpha'_0(X_m, X_{m+1})}{\alpha_0} \middle| X_{-j+2}, \mathbf{Y}_{-j+2,1} \right\} \\
&\quad - E_0 \left\{ \sum_{m=-j+2}^0 \frac{g'_0(Y_m|X_m)}{g_0} + \sum_{m=-j+2}^{-1} \frac{\alpha'_0(X_m, X_{m+1})}{\alpha_0} \middle| X_{-j+2}, \mathbf{Y}_{-j+2,0} \right\}
\end{aligned}$$

by the usual formula. Consider the first part of the m th term in the sum in (e),

$$\begin{aligned}
& U_{jm}^{(1)} \tag{f} \\
&\equiv E_0 \left\{ \frac{g'_0(Y_m|X_m)}{g_0} \middle| X_{-j+2}, \mathbf{Y}_{-j+2,1} \right\} - E \left\{ \frac{g'_0(Y_m|X_m)}{g_0} \middle| X_{-j+2}, \mathbf{Y}_{-j+2,0} \right\} \\
&= \sum_{a=1}^K \frac{g'_0(Y_m|a)}{g_0} \{P_0[X_m = a|X_{-j+2}, \mathbf{Y}_{-j+2,1}] - P_0[X_m = a|X_{-j+2}, \mathbf{Y}_{-j+2,0}]\}.
\end{aligned}$$

Note that, by the (backward) martingale convergence theorem, for fixed $m < 0$,

$$U_{jm}^{(1)} \xrightarrow{P_0} E_0 \left\{ \frac{g'_0(Y_m|X_m)}{g_0} \middle| Y_1, Y_0, \dots \right\} - E_0 \left\{ \frac{g'_0(Y_m|X_m)}{g_0} \middle| Y_0, Y_{-1}, \dots \right\} \tag{g}$$

as $j \rightarrow \infty$.

Note that,

$$\begin{aligned}
& P_0 \{X_m = a|X_{-j+2}, \mathbf{Y}_{-j+2,0}\} \tag{h} \\
&= \sum_b P_0 \{X_m = a|X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = b\} P_0 \{X_1 = b|X_{-j+2}, \mathbf{Y}_{-j+2,0}\}
\end{aligned}$$

and

$$\begin{aligned}
& P_0 \{X_m = a|X_{-j+2}, \mathbf{Y}_{-j+2,1}\} \tag{i} \\
&= \sum_c P_0 \{X_m = a|X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = c\} P_0 \{X_1 = c|X_{-j+2}, \mathbf{Y}_{-j+2,1}\}
\end{aligned}$$

so that,

$$\begin{aligned}
& \max_a |P_0 \{X_m = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}\} - P_0 \{X_m = a | X_{-j+2}, \mathbf{Y}_{-j+2,1}\}| \quad (\text{j}) \\
& \leq \max_{a,b,c} \left| P_0 \{X_m = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = b\} \right. \\
& \quad \left. - P_0 \{X_m = a | X_{-j+2}, \mathbf{Y}_{-j+2,0}, X_1 = c\} \right| \\
& = S(j, -m).
\end{aligned}$$

We conclude by lemma 3.17 that,

$$\begin{aligned}
|U_{jm}^{(1)}| & \leq 2\gamma^{-1}(\vartheta_0) \sum_{a=1}^K \left| \frac{g'_0}{g_0}(Y_m | a) \right| \prod_{k=m+1}^0 (1 - 2\mu_0(Y_k)) \quad (\text{k}) \\
& \leq 2\gamma^{-1}(\vartheta_0) K q_0(Y_m, M\delta_n) \exp \left(-2 \sum_{k=m+1}^0 \mu_0(Y_k) \right).
\end{aligned}$$

Now, by (k)

$$\begin{aligned}
& E_0 \left(\sum_{m=-j+2}^{-k} U_{jm}^{(1)} \right)^2 \quad (1) \\
& \leq 4\gamma^{-2}(\vartheta_0) K^2 \sum_{m_1=-j+2}^{-k} \sum_{m_2=-j+2}^{-k} E_0 \left\{ q_0(Y_{m_1}, M\delta_n) q_0(Y_{m_2}, M\delta_n) \right. \\
& \quad \left. \exp \left[-2 \left(\sum_{t=m_1}^0 \mu_0(Y_t) + \sum_{t=m_2}^0 \mu_0(Y_t) \right) \right] \right\}.
\end{aligned}$$

Applying the Hölder inequality to each term and using A3 we obtain

$$\begin{aligned}
& E_0 \left\{ \left(\sum_{m=-j+2}^{-k} U_{jm}^{(1)} \right)^2 \right\} \quad (\text{m}) \\
& \leq C_\epsilon \sum_{m_1} \sum_{m_2} E_0^{(1+\epsilon)^{-1}} \exp \left[-2(1+\epsilon) \left(\sum_{t=m_1}^0 \mu_0(Y_t) + \sum_{t=m_2}^0 \mu_0(Y_t) \right) \right].
\end{aligned}$$

But, if $m_1 \leq m_2$

$$E_0 \left\{ \exp \left[-2(1+\epsilon) \left(\sum_{t=m_1}^0 \mu_0(Y_t) + \sum_{t=m_2}^0 \mu_0(Y_t) \right) \right] \right\} \quad (\text{n})$$

$$\begin{aligned}
&= E_0 \left\{ \prod_{t=m_2}^0 E_0 \left(e^{-4(1+\epsilon)\mu_0(Y_t)} \middle| X_t \right) \prod_{t=m_1}^{m_2-1} E_0 \left(e^{-2(1+\epsilon)\mu_0(Y_t)} \middle| X_t \right) \right\} \\
&\leq \gamma_{4(1+\epsilon)}^{-m_2} \gamma_{2(1+\epsilon)}^{m_2-m_1} \\
&\leq \gamma_{2(1+\epsilon)}^{|m_1|}
\end{aligned}$$

where $\gamma_s = \max_a E_0(e^{-s\mu_0(Y_1)} | X_1 = a) < 1$ for all $s > 0$. Using the bound from (n) in (m) we obtain, for some $C_\epsilon < \infty$, $\gamma \equiv \gamma_{2(1+\epsilon)}$,

R27: Changed

$$\begin{aligned}
E_0 \left\{ \left(\sum_{m=-j+2}^{-k} U_{jm}^{(1)} \right)^2 \right\} &\leq 2C_\epsilon \sum_{m=k}^{j-2} m\gamma^{m(1+\epsilon)^{-1}} \quad (\text{o}) \\
&\leq 2C_\epsilon \gamma^{k(1+\epsilon)^{-1}} (1 - \gamma^{(1+\epsilon)^{-1}})^{-1}.
\end{aligned}$$

Thus for any $\delta > 0$ there exists $k = k(\delta)$ such that, for all $j > k + 2$,

$$E_0 \left\{ \left(\sum_{m=-j+2}^{-k} U_{jm}^{(1)} \right)^2 \right\} \leq \delta. \quad (\text{p})$$

A similar argument shows that for fixed k , some $C < \infty$, all j ,

$$E_0 \left(\sum_{m=-k}^0 U_{jm}^{(1)} \right)^4 \leq C. \quad (\text{q})$$

By a similar but easier argument if

$$\begin{aligned}
U_{jm}^{(2)} &= E_0 \left\{ \frac{\alpha'_0}{\alpha_0} (X_m, X_{m+1}) \middle| X_{-j+2}, \mathbf{Y}_{-j+2,1} \right\} \quad (\text{r}) \\
&\quad - E_0 \left\{ \frac{\alpha'_0}{\alpha_0} (X_m, X_{m+1}) \middle| X_{-j+2}, \mathbf{Y}_{-j+2,1} \right\}
\end{aligned}$$

Then

$$U_{jm}^{(2)} \xrightarrow{P_3} E_0 \left\{ \frac{\alpha'_0}{\alpha_0} (X_m, X_{m+1}) \middle| Y_1, Y_0, \dots \right\} - E_0 \left\{ \frac{\alpha'_0}{\alpha_0} (X_m, X_{m+1}) \middle| Y_0, Y_{-1}, \dots \right\} \quad (\text{s})$$

and (p) and (q) carry over as well. We conclude that (d) follows since in fact, by (g), (p)–(s),

$$U_j(X_{-j+2}, \mathbf{Y}_{-j+2,1}) \xrightarrow{L_2} W(Y_1, Y_0, \dots). \quad (\text{t})$$

The lemma follows. □

Proof of lemma 2.4: We begin with proving (2.12). In view of lemma 2.3 it is enough to show that for all τ ,

$$\text{Var}_0 \left(\frac{1}{n} \sum_{m=1}^N \tau^T E_0 \left\{ \nabla \nabla^T \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) | X_{mk+1} \right\} \tau^T \right) \rightarrow 0. \quad (\text{a})$$

But if we let $h_{k,m}(X_{mk+1})$ denote the m th summand in (a), lemma 3.7 and geometric ergodicity of the $\{X_j\}$ guarantees that the expression in (a) is bounded by,

$$C E_0 h_{k,1}^2(X_1) N n^{-2}. \quad (\text{b})$$

Also,

$$\begin{aligned} E_0 h_{k,1}^2(X_1) & \leq M^4 E_0 |\nabla \log \ell_0(\mathbf{Y}_1^{(k)} | X_1)|^4 \\ & \leq M^4 E_0 \left| \sum_{i=1}^k \nabla \log g_0(Y_i | X_i) + \sum_{i=1}^{k-1} \nabla \log \alpha_0(X_i, X_{i+1}) + \frac{\pi'_0}{\pi_0}(X_1) \right|^4 \\ & = O(k^2) \end{aligned} \quad (\text{c})$$

by invoking the formula of (e) of lemma 2.3 and lemma 3.7 again. Thus, $E_0 h_{k,1}^2(X_1) N n^{-2} = O(k n^{-1}) = o(1)$ and (a) and (2.12) follow. To prove (2.13) we take expectations and note that it is enough to show,

$$E_0 |\nabla \log \ell_0(\mathbf{Y}_1^{(k)} | X_1)|^4 = O(k^2). \quad (\text{d})$$

But this is just (c). Finally, (2.14) follows from,

$$\begin{aligned} P_0[\delta_n |\nabla \log \ell_0(\mathbf{Y}_1^{(k)} | X_1)| \geq \epsilon] & \leq n^{-2} \epsilon^{-4} E_0 |\nabla \log \ell_0(\mathbf{Y}_1^{(k)} | X_1)|^4 \\ & = O(k^2 n^{-2}). \end{aligned} \quad (\text{e})$$

□

Proof of lemma 2.5: By a standard identity valid under our conditions.

$$\left\| E_0 \left(\frac{\partial^2}{\partial \vartheta_a \partial \vartheta_b} \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \right) \right\| = -E_0(\nabla \nabla^T \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1})). \quad (\text{a})$$

Therefore, by lemma 2.3 and stationarity

$$\frac{1}{n} E_0 \sum_{m=1}^N \left\| \frac{\partial^2}{\partial \vartheta_a \partial \vartheta_b} \log \ell_0(\mathbf{Y}_m^{(k)} | X_{mk+1}) \right\| \rightarrow -I(\vartheta_0). \quad (\text{b})$$

Now use A5 and argue as in the proof of (2.12) to obtain the lemma.

□

4 References

- Ball, F. and Rice, J. (1992). Stochastic models for ion channels: introduction and bibliography. *Math. Biosciences* **112** 189-206.
- Baum, L. and Petrie, T. (1966). Statistical inference for probabilistic functions of finite state Markov chains. *Ann. Math. Statist.* **37** 1554-1563.
- Churchill, G. (1992). Hidden Markov chains and the analysis of genome structure. *Computers in Chemistry* **16** 107-115.
- Daley, D. and Vere Jones, D. (1989). *Introduction to the theory of point processes* Springer Verlag, New York.
- Frigessi, A. and Piccioni, M. (1990). Parameter estimation for 2 dimensional Ising fields corrupted by noise. *Stoch. Processes and their Applications* **34** 297-311.
- Furstenberg, H. and Kesten, H. (1960). Products of random matrices. *Ann. Math. Statist.* **31** 457-469.
- Götze, F. and Hipp, C. (1992). On the validity of Edgeworth expansions. Tech Report, University of Bielefeld.
- Hall, P. and Heyde, C. (1980). *Martingale limit theory and its applications*. Academic Press, New York.
- Hasminskii, R.Z. and Lazareva, B.V. (1992). Filtration procedures for jump Markov processes. *Ann. Statist.* **20** 2153-2164.
- Ibragimov, I. and Linnik, Y. (1971). *Independent and stationary sequences of random variables*. Walters Noordhoff, Groningen.
- Kingman, J.F.C (1976). Subadditive processes. In *Ecole d'Eté de Probabilitéés de Saint-Flour V-1976*, Ed: P.L. Hennequin, Lecture Notes in Mathematics 539, pp. 167-223, *Springer-Verlag*, Berlin.
- LeCam, L. (1956) On the asymptotic theory of estimation and testing hypotheses. *Third Berkeley Symp. Math. Statist. and Prob.* **1**, 129–156, Univ. of California Press, Berkeley.

- LeCam, L. (1986). *Asymptotic methods in statistical decision theory*. Springer-Verlag, New York. R28: Changed
- LeCam, L. and Yang, G. (1990). *Asymptotics in statistics. Some Basic Concepts*. Springer Verlag, New York.
- Leroux, B. (1989). Maximum likelihood estimation for mixture distributions and hidden Markov models. *Thesis Univ. of British Columbia*.
- Leroux, B. (1992). Maximum likelihood estimation for hidden Markov models. *Stoch. Processes and their Applications* **40** 127-143.
- Petrie, T. (1969). Probabilistic functions of finite state Markov chains. *Ann. Math. Statist.* **40** 97-115.
- Priestley, M.B. (1988). *Nonlinear and nonstationary time series* Academic Press, London.
- Rabiner, L.R. (1989). A tutorial on hidden Markov models and selected applications in speech recognition. *Proc. IEEE* **77** 257-284.
- Rydén, T. (1994a) Consistent and asymptotically normal parameter estimates for hidden Markov models. *Ann. Statist.*, to appear R29: Changed
- Rydén, T. (1994b). Parameter estimation for Markov modulated Poisson Processes. *Stochastic Models* **10**, 795–829.
- Rydén, T. (1995) Estimating the order of hidden Markov models, *Statistics*, to appear.
- Titterton, D. M. (1991) Estimation of parameters in hidden Markov models. *Proc. Roy. Soc. Lond.* **A 337**, 407–428.
- Tong, H. (1991). *Nonlinear time series* Oxford University Press.